




BALANCER

PELLET

50 LBS
(22.7 KG)



IDEAL FOR:

Overweight horses, broodmares, growing horses, metabolic horses



FEATURES:

Whole grain & molasses free, low starch & sugar, low feeding rate



HELPS SUPPORT:

Balancing hay & pasture, feeding grains below recommended feeding rate

While originally formulated for growing horses and broodmares, Triple Crown Balancer is a highly fortified, low sugar and starch pelleted supplement, ideal for easy keepers and horses with metabolic disease and insulin resistance. This can be used to balance pasture and/or hay and improve nutrition in diets when feeding below the recommended feeding rate and/or mixing with whole grains.

PROTEIN 30% • FAT 3% • FIBER 5% • 1,266 KCAL/LB

GUARANTEED ANALYSIS

Crude Protein (min.)	30.00%
Lysine (min.)	2.00%
Methionine (min.)	0.62%
Threonine (min.)	1.35%
Tryptophan (min.)	0.44%
Leucine (min.)	2.25%
Crude Fat (min.)	3.00%
Crude Fiber (max.)	5.00%
ADF (max.)	7.00%
NDF (max.)	13.00%
Calcium (min.)	4.00%
Calcium (max.)	5.00%
Phosphorus (min.)	1.50%
Magnesium (min.)	2.40%
Potassium (min.)	1.30%
Sodium (min.)	0.50%
Sodium (max.)	1.00%
Salt (min.)	0.95%
Salt (max.)	1.50%
Omega-3 Fatty Acids (min.)	0.40%
Omega-6 Fatty Acids (min.)	1.10%
Selenium (min.)	3.40 ppm
Selenium (max.)	4.40 ppm
Zinc (min.)	1,000 ppm
Manganese (min.)	575 ppm
Cobalt (min.)	2.10 ppm
Copper (min.)	300 ppm
Iron (min.)	500 ppm
Vitamin A (min.)	36,000 IU/lb.
Vitamin D (min.)	6,000 IU/lb.
Vitamin E (min.)	1,000 IU/lb.
Riboflavin-B2 (min.)	16.00 mg/lb.
Thiamin-B1 (min.)	42.00 mg/lb.
Ascorbic Acid (min.)	275 mg/lb.
Biotin (min.)	1.40 mg/lb.
Total Direct Fed Microbials (min.)	16.85 billion CFU/lb.
Lactobacillus Acidophilus Fermentation Product (min.)	4.54 billion CFU/lb.
Enterococcus Faecium Fermentation Product (min.)	4.54 billion CFU/lb.
Bacillus Subtilis Dried Fermentation Product (min.)	125 million CFU/lb.
Saccharomyces Cerevisiae Yeast Culture (min.)	7.65 billion CFU/lb.
Cellulase ¹ (Trichoderma Longibrachiatum Fermentation Extract) (min.)	3.20 million CFU/lb.
Protease ² (Bacillus Subtilis Fermentation Extract) (min.)	7150 HUT/lb.

INGREDIENTS

(Free from Restricted Ruminant Protein Products per Title 21, CFR 589.2000/2001) Dehulled Soybean Meal, Wheat Middlings, Ground Flaxseed, Stabilized Rice Bran, Calcium Carbonate, Monocalcium Phosphate, Dicalcium Phosphate, Magnesium Oxide, Yeast Extract, Yeast Culture, Hydrolyzed Yeast, Dried Enterococcus Faecium Fermentation Product, Dried Lactobacillus Acidophilus Fermentation Product, Dried Bacillus Subtilis Fermentation Product, Dried Trichoderma Longibrachiatum Fermentation Extract, Dried Bacillus Subtilis Fermentation Extract, Zinc Hydroxy Chloride, Manganese Hydroxy Chloride, Copper Hydroxy Chloride, Vegetable Oil, Butyric Acid, Zinc Oxide, Peppermint Essential Oil, Kelp Meal, Lecithin, Magnesium Proteinate, Selenium Yeast, L-Lysine, DL-methionine, L-Threonine, L-Leucine, Soybean Oil, Cane Molasses, Salt, Sodium Bicarbonate, Fenugreek Seed, Anise, Ascorbic Acid (Source of Vitamin C), Niacin Supplement, Biotin, Vitamin A Supplement, Vitamin E Supplement, Thiamine Mononitrate, Beta Carotene, Calcium Pantothenate, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B12 Supplement, Vitamin D3 Supplement, Choline Chloride, Menadione Sodium Bisulfite Complex (Source of Vitamin K Activity), Folic Acid, Hydrated Sodium Calcium Aluminosilicate, Brewers Dried Yeast, Manganese Sulfate, Zinc Sulfate, Copper Sulfate, Cobalt Proteinate, Iron Amino Acid Complex, Ethylenediamine Dihydroiodide, (Propionic Acid, Sodium Benzoate, Potassium Sorbate (Preservatives)).

FEEDING DIRECTIONS

TO BALANCE PASTURE AND/OR HAY³

Ponies: 500 – 700 lb.	0.5 - 0.75 lb. per day
Weanlings: 400 – 600 lb.	0.5 - 0.75 lb. per day
Yearlings: 700 – 800 lb.	1 lb. per day
Horses: 800 – 1,100 lb.	1 - 1.5 lb. per day
Broodmares and Breeding Stallions	1.5 - 2 lb. per day
Horses in moderate-to-high levels of activity	1.5 - 2 lb. per day

TO BALANCE DIETS WHEN FEEDING BELOW RECOMMENDED FEEDING RATE

Ponies: 500 – 700 lb.	0.25 - 0.5 lb. per day
Weanlings: 400 – 600 lb.	0.5 - 0.75 lb. per day
Yearlings: 700 – 800 lb.	1 lb. per day
Horses: 800 – 1,100 lb.	0.5 - 1 lb. per day
Broodmares and Breeding Stallions	1 lb. per day
Horses in moderate-to-high levels of activity	1 lb. per day

TO BALANCE GRAINS⁴

Ponies: 500 – 700 lb.	0.5 - 0.75 lb. per day
Horses: 800 – 1,100 lb.	1 - 1.5 lb. per day
Broodmares and Breeding Stallions	1.5 - 2 lb. per day
Horses in moderate-to-high levels of activity	1.5 - 2 lb. per day

OTHER CONSIDERATIONS

The above feeding recommendations assume a horse is consuming a minimum of 1.5 to 2% of its body weight in hay or pasture.

Provide salt and plenty of fresh, clean water at all times. Keep product fresh in cool, dry storage. Examine product daily. DO NOT use if old, moldy or contaminated with insects.

1. One cellulase unit (CU) is defined as the amount of activity that will produce a relative fluidity change of 1 (determined with a calibrated viscometer) in 5 minutes in a defined carboxymethylcellulose substrate at pH 4.5 and 40°C.

2. One hemoglobin unit (HUT) of proteolytic (protease) activity is defined as that amount of enzyme that produces, in one minute under the specified conditions, a hydrolysate whose absorbance at 275nm is the same as that of a solution containing 1.10g per mL of tyrosine in 0.006N hydrochloric acid. Estimated values determined by Equi-Analytical, Ithaca, NY.

3. Assuming a horse is consuming a minimum of 1.5 to 2% of its body weight in hay or pasture.

4. The feeding recommendations provide for appropriate vitamin/mineral fortification according to body weight. For desired body condition, adjust grain inclusion (oats, corn, barley) to provide additional calories.