

GOATGROW
GOAT GROWER DEVELOPER
PELLETS

Formulated for Goats

MEDICATED

For the prevention of coccidiosis caused by *Eimeria crandalis*, *Eimeria christenseni* and *Eimeria ninakohlyakimovae*.

ACTIVE DRUG INGREDIENT

Monensin (as monensin sodium)....20 g/ton

GUARANTEED ANALYSIS

Crude Protein, min.....16.0%
(Contains not more than 1.0% protein from non-protein nitrogen.)
Crude Fat, min.....2.5%
Crude Fiber, max.....16.0%
Calcium, min...0.9%....max....1.6%
Phosphorus, min.....0.3%
Salt, min....0.8%.....max...1.5%
Copper, min...51ppm....max..52ppm
Vitamin A, min.....9.0 KIU/lb
Selenium, min.....0.27 ppm

INGREDIENTS

Roughage Products, Grain Products, Processed Grain By-Products, Plant Protein Products, Cane Molasses, Vegetable Fat Product (Feed Grade), Propylene Glycol, Lecithin, Phosphoric Acid, Sulfuric Acid, Caramel Flavoring, Propionic Acid, Acetic Acid, and Benzoic Acid (as preservatives), Calcium Carbonate, Salt, Ammonium Chloride, Spray Dried Lignin Sulfonate, Monocalcium Phosphate, Dicalcium Phosphate, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Calcium Iodate, Cobalt Carbonate, Sodium Bentonite, and Mineral Oil, Potassium Chloride, Magnesium Oxide, Sodium Selenite, Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E Supplement, Zinc Methionine.

DIRECTIONS

Feed continuously as the sole ration with good quality forage & fresh, clean water. Quantity will vary with the bodyweight conditions of the goats being fed.

Notice: Type C goat feeds shall bear an expiration date of 30 days after date of manufacture.

Caution: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by equines has been fatal. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal in cattle and could be fatal in goats. Monensin medicated feed is intended for use in cattle or goats. Consumption by unapproved species may result in toxic reactions. Do not feed to lactating goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the level of Monensin recommended in the feeding direction, as reduced average daily gains may result.

Manufactured By:
AUGUSTA COOPERATIVE FARM BUREAU, INC.
General Offices Staunton, Virginia 24401
NET WEIGHT 50 lbs. (22.68 kg)
MEDICATED